

Wind Stress Functions for Blowing Snow Initiation and Transport, in Vegetated Grids

KEMP I. SIMON¹, PETER A. TAYLOR¹, AND MARK GORDON¹

EXTENDED ABSTRACT

Adequate representation of snow processes in land surface schemes would improve General Circulation Model (GCM) simulations. The Canadian Land Surface Scheme (CLASS) and other land surface models lack blowing snow codes. A blowing snow sublimation parameterization recently formulated from simplifying assumptions and the PIEKTUK model results, omits micro-scale details, making it suitable for incorporation into CLASS. The parameterization is specific to non-vegetated surfaces and its adaptation for use in vegetated grids is being targeted here. The transport formula for PIEKTUK and the Prairie Blowing Snow Model (PBSM) are similar except for vegetation roughness represented only in the latter. This similarity provides the opportunity to formulate vegetated transport rates on the basis of PBSM results and to fit the formula to the new blowing snow sublimation parameterization to generate vegetated sublimation rates.

In this approach, a wind stress function influences threshold, or otherwise effective, velocities for testing blowing snow initiation at various vegetation exposures. For strong-wind situations a wind dependent stress function, formulated from the non-vegetated to vegetated wind stress ratio, generates transport rates comparable to those of the PBSM. These transport rates increase as a power function of the wind speed and are higher for smoother surfaces. With weaker winds blowing snow sublimation is limited to the upper canopy, hence the assessed depth of penetration of the effective wind, control the vegetated-grid blowing snow transport and sublimation rates. This depth derives from an empirical formula representing the relative within-canopy to canopy height wind stress, on the basis of previous measurements by others.

The resulting wind velocities of blowing snow initiation are temperature and roughness influenced. Together with the parameterized non-vegetated sublimation rates, these threshold velocity and transport functions allow us to represent blowing snow sublimation from vegetated grids via the Canadian Land Surface Scheme. As an application we note that parameterization of blowing snow sublimation with partial vegetation cover and roughness averaging, gives a slightly improved representation of the Resolute snow depth profile for 1966–1971. Our simple parameterization can be improved by the incorporation of snow ageing in the threshold wind velocity formula as for the improved PBSM model. We also suggest the modeling of snow densification due to wind packing, in CLASS, and the representation of its influence on the threshold velocity.

Key Words: Blowing Snow, Land Surface Schemes, Transport, Sublimation, Vegetation.

REFERENCES

- Cess RD. et al, 1991. Interpretation of Snow-Climate Feedbacks as Produced by 17 General Circulation Models. *Science* **253**: 888–892.
- Déry SJ, Taylor PA, Xiao J. 1998. Thermodynamic effects of Sublimating Blowing Snow in the Atmospheric Boundary Layer. *Boundary-Layer Meteorology* **93**: 237–251.
- Garratt JR. 1992. *The Atmospheric Boundary Layer*. Cambridge University Press, 316 pp.
- Gordon M, Simon K, Taylor P. 2005. A Parameterization of Blowing Snow Sublimation within the Canadian Land Surface Scheme. *Atmosphere-Ocean*, submitted.
- Kind RJ. 1981. Snow Drifting. In Gray DM, Male DH. (eds.). *Hand-book of snow—Principles, Processes, Management & Use*. Pergamon press, 383–358.
- Li L, Pomeroy JW. 1997. Estimates of Threshold Wind Speeds for Snow Transport Using Meteorological Data. *Journal of Applied Meteorology* **36**: 205–213.
- Mason PJ. 1988. The Formulation of Areally-Averaged Roughness Lengths. *Quarterly Journal of the Royal Meteorological Society* **114**: 399–420.
- Maykut GA, Church PE. 1973. Radiation and Climate of Barrow, Alaska, 1962–66. *Journal of Applied Meteorology* **12**: 620–628.
- Pomeroy JW. 1989. A Process-Based Model of Snow Drifting. *Annals of Glaciology* **13**: 237–240.
- Pomeroy JW. 1988. *Wind Transport of Snow*. PhD Thesis, Division of Hydrology, University of Saskatchewan, Saskatoon, 226 pp.
- Pomeroy JW, Li L. 2000. Prairie and Arctic Areal Snow Cover Mass Balance Using a Blowing Snow Model. *Journal of Geophysical Research* **105**: D21, 26619–26634.
- Pomeroy JW, Gray DM. 1995. *Snowcover Accumulation, Relocation and Management*. National Hydrological Research Institute, Environment Canada, Saskatoon. Sci. Rep. **7**: 144 pp.
- Pomeroy JW, Gray DM, Landine PG. 1993. The Prairie Blowing Snow Model; Characteristics, Validation, Operation. *Journal of Hydrology* **144**: 165–192.
- Pomeroy JW et al, 1998. An Evaluation of Snow Accumulation and Ablation Processes for Land Surface Modeling. *Hydrological Processes* **12**: 2339–2367.
- Raupach NR. 1988. Canopy Transport Processes. In Steffan WL, Denmead OT. Eds. *Flow and Transport in the Natural Environment: Advances and Applications*. Springer-Verlag, 95–127.
- Schmidt RA. 1991. Sublimation of snow intercepted by an artificial conifer. *Agricultural and Forest Meteorology* **54**: 1–27.
- Schmidt RA. 1982. Vertical Profiles of Wind Speed, Snow Concentrations and Humidity in Blowing Snow. *Boundary-Layer Meteorology* **23**: 223–246.
- Thorpe AD, Mason BJ. 1966. The Evaporation of Ice Spheres and Ice Crystals. *British Journal of Applied Physics* **17**: 541–548.
- Verseghy Diana L. 1991. CLASS—A Canadian Land Surface Scheme for GCMs I, Soil Model. *International Journal of Climatology* **11**: 111–133.
- Verseghy DL, McFarlane NA, Lazare M. 1993. CLASS—A Canadian Land Surface Scheme for GCMs, II. Vegetation Model and Coupled Runs. *International Journal of Climatology* **13**: 347–370.
- Xiao J. 2001. *A model of the Transport and Sublimation of Blowing Snow in the Atmospheric Boundary Layer*. PHd Thesis, York University, Toronto, 227 pp.